

Brief Articles

Design and Synthesis of Celecoxib and Rofecoxib Analogues as Selective Cyclooxygenase-2 (COX-2) Inhibitors: Replacement of Sulfonamide and Methylsulfonyl Pharmacophores by an Azido Bioisostere

Amgad G. Habeeb, P. N. Praveen Rao, and Edward E. Knaus*

Faculty of Pharmacy and Pharmaceutical Sciences, University of Alberta, Edmonton, Alberta, Canada

Received April 3, 2001

Celecoxib (**13**) and rofecoxib (**17**) analogues, in which the respective SO_2NH_2 and SO_2Me hydrogen-bonding pharmacophores were replaced by a dipolar azido bioisosteric substituent, were investigated. Molecular modeling (docking) studies showed that the azido substituent of these two analogues (**13**, **17**) was inserted deep into the secondary pocket of the human COX-2 binding site where it undergoes electrostatic interaction with Arg⁵¹³. The azido analogue of rofecoxib (**17**), the most potent and selective inhibitor of COX-2 (COX-1 IC_{50} = 159.7 μM ; COX-2 IC_{50} = 0.196 μM ; COX-2 selectivity index = 812), exhibited good oral antiinflammatory and analgesic activities.

Introduction

Many selective COX-2 inhibitors belong to a tricyclic group of compounds with a central ring possessing a diaryl stilbene-like structure with a sulfonyl (SO_2) group at the para position of one of the aryl rings, such as DuP-697 (**1**),¹ celecoxib (Celebrex) (**2**),² SC-588 (**3**),² rofecoxib (Vioxx) (**4**),³ 3-(4-methylsulfonylphenyl)-4-phenyl-3-trifluoromethylisoxazole (**5**),⁴ and 2,3-dimethyl-5-(4-methylsulfonylphenyl)-4-phenyl-4-isoxazoline (**6**),⁵ as illustrated in Figure 1. The SO_2Me and SO_2NH_2 pharmacophores are believed to induce COX-2 selectivity by insertion into the secondary pocket of COX-2 which is absent in COX-1. The secondary pocket present in COX-2 has been attributed to the presence of isoleucine (Ile⁵²³) in COX-1 relative to the smaller valine (Val⁵²³) in COX-2.⁶ Replacement of histidine (His⁵¹³) in COX-1 by arginine (Arg⁵¹³) in COX-2 has been reported to play a key role in the hydrogen-bond network of the COX active site. Histidine (His⁹⁰), glutamine (Gln¹⁹²), and tyrosine (Tyr³⁵⁵) control the access of ligands into the secondary pocket.⁷ The interaction of Arg⁵¹³ with the bound ligand has been reported to be a requirement for the time-dependent inhibition of COX-2.⁸ The presence of the Arg⁵¹³ residue, to our knowledge, has not been exploited for the design of selective COX-2 inhibitors. Accordingly, we now describe the design, synthesis, cyclooxygenase inhibitory, analgesic and antiinflammatory activities, and some molecular modeling studies for the pyrazole regioisomers **10** and **13** and the furanone **17** that possess an azido group in place of the SO_2NH_2 and SO_2Me pharmacophores present in celecoxib and rofecoxib, respectively.





Figure 1. Representative examples of selective COX-2 inhibitors having a central five-membered heterocyclic ring.

Chemistry

Reaction of 4-nitrophenylhydrazine with 1-(4-methylphenyl)-4,4,4-trifluorobutane-1,3-dione² in EtOH afforded the cyclic pyrazoline-5-ol **7** which eliminated a molecule of water upon treatment with HOAc at reflux temperature to yield the pyrazole **8** (see Scheme 1). Reduction of the nitro group in the pyrazole **8** with hydrazine hydrate and 10% Pd/C, using a method reported by Penning et al.,² yielded the corresponding amino product **9**. Diazotization of **9**, and treatment of the diazonium salt with NaN_3 , afforded the 1,3-regioisomer of celecoxib having an azido substituent in place of the SO_2NH_2 pharmacophore. In contrast, the 1,5-regioisomer was prepared by condensation of 4-nitro-

* Address correspondence to Dr. Edward E. Knaus, Faculty of Pharmacy and Pharmaceutical Sciences, University of Alberta, Edmonton, Alberta, Canada T6G 2N8. Tel: 780-492-5993. Fax: 780-492-1217. E-mail: eknaus@pharmacy.ualberta.ca.

Scheme 1^a


^a Reagents and conditions: (a) EtOH, reflux, 20 h; (b) HOAc, reflux, 2 h; (c) H₂NNH₂·xH₂O, 10% Pd/C, reflux, 45 min; (d) NaNO₂/HCl and then NaN₃, 0–5 °C, 45 min; (e) EtOH, HCl, reflux, 20 h.

phenylhydrazine with 1-(4-methylphenyl)-4,4,4-trifluorobutane-1,3-dione in EtOH under acidic reaction conditions to yield the pyrazole **11** (76%). This reaction, performed under acidic reaction conditions, provided a superior yield relative to related reactions carried out under neutral reaction conditions.² The azido analogue of celecoxib **13** was prepared starting from the pyrazole **11** using the same reaction sequence used for the elaboration of the nitro compound **10** to the azido product **11**.

The azido analogue **17** of rofecoxib, where MeSO₂ is replaced by N₃, was prepared starting with the bromination of 4-azidoacetophenone **14** using Br₂ according to a reported method.¹⁰ The subsequent reaction of the bromo compound **15** with phenylacetic acid in the presence of (Et)₃N gave 4-(4-azidophenyl)-3-phenyl-2(5H)furanone (**17**) that was formed via the intermediate ester **16** as illustrated in Scheme 2.

Results and Discussion

Celecoxib and rofecoxib analogues, having an azido group in place of the respective SO₂NH₂ and SO₂Me pharmacophores, were investigated to determine whether the azido substituent is a suitable bioisostere with respect to selective COX-2 inhibition, and AI and analgesic activities. Structure–activity studies for the tricyclic class of selective COX-2 inhibitors have shown that a SO₂Me or SO₂NH₂ substituent at the para position of one aryl ring usually confers optimal COX-2 inhibitory potency.¹¹ In the 1,2-diarylcyclopentene class

Scheme 2^a

^a Reagents and conditions: (a) Br₂, CHCl₃, 25 °C, 2 h; (b) PhCH₂COOH, Et₃N, CH₃CN, 25 °C, 1 h; (c) Et₃N, CH₃CN, reflux, 8 h.




Figure 2. Azide resonance hybrid structures.




Figure 3. Docking the pyrazole **13** (ball-and-stick) in the active site of human COX-2 (line and stick) ($E_{\text{intermolecular}} = -46.49$ kcal/mol). The C-atom of the CF₃ substituent is 10.26 Å from the phenolic OH of Tyr³⁵⁵, but removed from Ser⁵³⁰ (OH) by 7.18 Å. The terminal N-atom of the azido substituent is about 4.52 Å inside the entrance to the secondary pocket (Val⁵²³). The center of the N-1 phenyl ring is about 3.95 Å from the entrance to the secondary pocket (Val⁵²³).


of compounds, replacement of SO₂Me by SO₂CF₃, COMe, PO(OH)Me, CO₂H, PO(OH)₂, or SO(=NH)Me abolished COX-2 inhibitory activity.¹² A similar replacement of SO₂Me by NO₂, which can dispose a pair of oxygen atoms such as SO₂, in the 1,5-diarylpiprazole class also abolished COX-2 inhibitory activity.²

The azido substituent is particularly attractive since it has the potential to undergo electrostatic (ion–ion) binding interactions with amino acid residues, particularly Arg⁵¹³, lining the secondary pocket of COX-2. Covalent azides can be viewed as resonant hybrids between structures **A**, **B**, and **C** (see Figure 2).¹³ Pauling rejected **C** as a major contributor based on the *adjacent charge rule*.¹⁴ The remaining hybrids **A** and **B** predict a 2.5 bond order for the N₂–N₃ bond and a 1.5 bond order for the N₁–N₂ bond (see **D**, Figure 2). This prediction was in very good agreement with a structure determination of methyl azide (**D**) where the bond

Table 1. Antiinflammatory and Analgesic Activities, in Vitro COX-1 and COX-2 Inhibition Data, and Molecular Volumes of 1-(4-Azidophenyl)-3-(4-methylphenyl)-5-trifluoromethylpyrazole (**10**), 1-(4-Azidophenyl)-5-(4-methylphenyl)-3-trifluoromethylpyrazole (**13**), and 4-(4-Azidophenyl)-3-phenyl-2(5*H*)furanone (**17**)


| compd | AI activity ^a | | analgesic activity ^b | | vol. (Å ³) ^c | IC ₅₀ , μM ^d | | selectivity index (COX-1/COX-2) |
|-----------|--------------------------|-------------------------|---------------------------------|------------------------|-------------------------------------|------------------------------------|-------------------|---------------------------------|
| | % inhibition at 3 h | % inhibition at 5 h | % inhibition at 30 min | % inhibition at 60 min | | COX-1 | COX-2 | |
| 10 | | | | | 278.3 | 9.88 | 2.63 | 3.74 |
| 13 | 46.6 ± 4.4 | 16.9 ± 2.7 | 60.9 ± 9.4 | 63.1 ± 1.2 | 279.4 | >100 | 1.55 | 64.55 |
| 17 | 42.9 ± 1.0 | 27.5 ± 4.6 | 46.7 ± 1.3 | 60.6 ± 1.6 | 242.1 | 159.72 | 0.196 | 812.4 |
| celecoxib | 79.9 ± 1.9 ^e | 58.2 ± 1.8 ^f | 31.7 ± 9.6 | 62.0 ± 7.3 | 298.4 | 22.9 | 0.0507 | 404 |
| rofecoxib | | | | | 262.5 | 26.0 ^g | 0.34 ^g | 76.5 |

^a Inhibitory activity on carrageenan-induced rat paw edema; the result is the mean value ± SEM using four animals following a 50 mg/kg oral dose of the test compound. ^b Inhibitory activity in the rat 4% NaCl-induced abdominal constriction assay; the result is the mean value ± SEM using four animals following a 50 mg/kg ip dose of the test compound. ^c The volume of the molecule, after minimization using the MM3 force field, was calculated using the Alchemy 2000 program. ^d The result (IC₅₀, μM) is the mean of two determinations. ^e ID₅₀ = 10.8 mg/kg po dose. ^f ID₅₀ = 40.8 mg/kg po dose. ^g Data taken from the literature for inhibition of purified human recombinant COX-1 and COX-2.¹⁹

**Figure 4.** Docking the pyrazole (**10**) (ball-and-stick) in the active site of human COX-2 (line and stick) ($E_{\text{intermolecular}} = -34.12$ kcal/mol). The C-atom of the CF₃ substituent is 12.25 Å from the phenolic OH of Tyr³⁵⁵, but removed from the Ser⁵³⁰ (OH) by 8.56 Å. The terminal N-atom of the azido substituent is about 10.30 Å outside the entrance to the secondary pocket (Val⁵²³). The center of the N-1 phenyl ring is about 6.46 Å outside the entrance to the secondary pocket (Val⁵²³).

lengths from electron diffraction studies¹⁵ were as follows: N₂–N₃ = 1.12 Å, N₁–N₂ = 1.24 Å, C–N₁ = 1.47 Å, and the C–N₁–N₂ bond angle was 120°. The linear configuration of the azido group is in agreement with the sp³ hybridization indicated by the lack of nonbonded electron pairs on N₂.¹³ The azido group is slightly smaller in size [MR (molar refractivity) = 10.20] than a SO₂Me (MR = 13.49) or SO₂NH₂ (MR = 12.28) substituent, but more lipophilic (π = 0.46) relative to the more polar SO₂Me (π = –1.63) and SO₂NH₂ (π = –1.82) substituents¹⁶ which have the potential to improve absorption and provide a more rapid onset of action.¹¹

Docking 1-(4-azidophenyl)-5-(4-methylphenyl)-3-trifluoromethylpyrazole (**13**) in the active site of human COX-2 (1CX2 PDB file), showed that the terminal N-atom of the azido group was inserted into the secondary COX-2 pocket about 4.52 Å from Val⁵²³, and about 3.15 Å from the center of the guanidino group of Arg⁵¹³ (see Figure 3). This orientation of the pyrazole **13** within the COX-2 active site provides an intermolecular energy between the enzyme and pyrazole **13** of about –46.19 kcal/mol, where the electrostatic component accounts for about 12% of this total energy. In comparison, the celecoxib (**2**) docked complex showed an intermolecular energy between the enzyme and celecoxib of about

**Figure 5.** Docking the furanone **17** (ball-and-stick) in the active site of human COX-2 (line and stick) ($E_{\text{intermolecular}} = -49.6$ kcal/mol). The O-atom of the CO substituent is 13.51 Å from the phenolic OH of Tyr³⁵⁵, but removed from Ser⁵³⁰ (OH) by 4.03 Å. The terminal N-atom of the azido substituent is about 4.52 Å inside the entrance to the secondary pocket (Val⁵²³). The center of the C-4 phenyl ring is about 4.11 Å from the entrance to the secondary pocket (Val⁵²³).

–45.16 kcal/mol where 2.6% was due to an electrostatic component. This difference is likely due to a greater electrostatic interaction between the dipolar azido group in pyrazole **13** and the charged guanidino moiety of Arg⁵¹³ in the secondary COX-2 pocket. Similar docking of the 1-(4-azidophenyl)-3-(4-methylphenyl)-5-trifluoromethylpyrazole regioisomer (**10**) in the active site of human COX-2 showed that the azido substituent did not insert into the secondary pocket since the ligand **10** extended parallel to the longitudinal axis of the hydrophobic primary COX-2 channel (cavity), in a manner characteristically observed for nonselective COX-2 inhibitors¹⁷ as illustrated in Figure 4.

These molecular modeling studies correlate well with in vitro enzyme inhibition data. In this regard, the 1,5-pyrazole regioisomer **13** showed selective inhibition of COX-2 [COX-1 IC₅₀ > 100 μM; COX-2 IC₅₀ = 1.5 μM; selectivity index (SI) ≈ 64], whereas the 1,3-regioisomer **10** showed a modest COX-2 selectivity ≈ 4. These results are similar to those described for other studies^{2,18} utilizing compounds not having a 1,2-diarylstilbene-like structure. Docking the rofecoxib analogue **17**, in which

the SO₂Me moiety was replaced by an azido substituent, in the active site of human COX-2 showed a similar interaction between the azido group and the COX-2 secondary pocket amino acid residues similar to that observed for the pyrazole **13** (see Figure 5). The intermolecular energy between the ligand **17** and the enzyme was -49.6 kcal/mol with the electrostatic component comprising 5.8% of the total energy. In contrast, the rofecoxib (**4**) docked complex showed an intermolecular energy of -42.16 kcal/mol where only 1.2% was due to an electrostatic component. The higher electrostatic component for the azido compound **17** (5.8%), relative to that for rofecoxib **4** (1.2%), is attributed to the fact that the MeSO₂ moiety present in rofecoxib undergoes H-bonding to the imidazole NH of His⁹⁰ in the secondary pocket. In contrast, the azido moiety in **17** undergoes an electrostatic interaction with Arg⁵¹³ in the secondary COX-2 pocket. The lower contribution of the electrostatic energy to the total intermolecular energy in the case of the furanone **17**, relative to the pyrazole **13**, can be attributed to the observed hydrogen bonding interaction between the O-atom of the C=O in the furanone structure and residues lining the primary COX-2 channel, particularly Arg¹²⁰. The azido analogue of rofecoxib **17** exhibited a potent and selective inhibition of COX-2 (COX-2 IC₅₀ = 0.196 μM; COX-1 IC₅₀ = 159.7 μM; SI ≈ 812). The molecular volumes of the selective COX-2 inhibitors **13** (279.4 Å³) and **17** (242.1 Å³) are moderately smaller than that for the selective COX-2 inhibitors celecoxib (298.4 Å³) and rofecoxib (262.5 Å³).

The azido analogues of celecoxib **13** and rofecoxib **17** exhibited good AI and analgesic activities (see Table 1).

Conclusions

In conclusion, the dipolar azido group is a bioisostere of the SO₂NH₂ and SO₂Me hydrogen-bonding pharmacophores present in many selective COX-2 inhibitors, and the azido analogues **13** and **17** may be useful biochemical agents for photoaffinity labeling of the COX-2 enzyme.

Experimental Section

Cyclooxygenase Inhibition Studies. All compounds described herein were tested for their ability to inhibit COX-1 and COX-2 using a COX-(ovine) inhibitor screening kit (catalog no. 560101, Cayman Chemical, Ann Arbor, MI) using the method previously reported.⁴

Antiinflammatory Assay. The test compounds were evaluated using the in vivo rat carrageenan-induced foot paw edema model reported previously.²⁰

Analgesic Assay. Analgesic activity was determined using the 4% sodium chloride-induced writhing (abdominal constriction) assay as described previously.²¹

Acknowledgment. We are grateful to the Canadian Institutes of Health Research (Grant MOP-14712) for financial support of this research and to C.-A. McEwen for technical assistance in performing the antiinflammatory and analgesic assays.

Supporting Information Available: Experimental procedures for the preparation of compounds **7–13** (Scheme 1) and **15–17** (Scheme 2) and their IR and NMR (¹H, ¹³C, ¹⁹F) spectroscopic data. This material is available free of charge on the Internet at <http://pubs.acs.org>.

References

- Gans, K. R.; Galbraith, W.; Roman, R. J.; Haber, S. B.; Kerr, J. S.; Schmidt, W. K.; Smith, C.; Hewes, W. E.; Ackerman, N. R. Antiinflammatory and safety profile of DuP-697, a novel orally effective prostaglandin synthesis inhibitor. *J. Pharmacol. Exp. Ther.* **1990**, *254*, 180–187.
- Penning, T.; et al. Synthesis and biological evaluation of the 1,5-diarylpyrazole class of cyclooxygenase-2 inhibitors: Identification of 4-[5-(4-methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzenesulfonamide (SC-58635, Celecoxib). *J. Med. Chem.* **1997**, *40*, 1347–1365.
- Desmond, R.; Dolling, U.; Marcune, B.; Tillyer, R.; Tschaen, D. Process for making phenylheterocycles useful as COX-2 inhibitors. *World Patent* WO96/08482, 21 March 1996; *Chem. Abstr.* **1996**, *125*, P 86474.
- Habeeb, A. G.; Praveen Rao, P. N.; Knaus, E. E. Design and syntheses of 4,5-diarylloxazoles: Novel inhibitors of cyclooxygenase-2 (COX-2) with analgesic and antiinflammatory activity. *Drug Devel. Res.* **2000**, *51*, 273–286.
- Habeeb, A. G.; Praveen Rao, P. N.; Knaus, E. E. Design and Synthesis of 4,5-diphenyl-4-isoxazolines: Novel inhibitors of Cyclooxygenase-2 (COX-2) with an analgesic and antiinflammatory activity. *J. Med. Chem.* In press.
- Luong, C.; Miller, A.; Barnett, J.; Chow, J.; Ramesha, C.; Browner, M. F. Flexibility of the NSAID binding site in the structure of human cyclooxygenase-2. *Nat. Struct. Biol.* **1996**, *3*, 927–933.
- Llorens, O.; Perez, J. L.; Palomer, A.; Mauleon, D. Structural basis for dynamic mechanism of ligand binding to COX. *Biorg. Med. Chem. Lett.* **1999**, *9*, 2779–2784.
- Garavito, R. M.; DeWitt, D. L. The Cyclooxygenase isoforms: Structural insights into the conversion of arachidonic acid to prostaglandins. *Biochim. Biophys. Acta* **1999**, *1441*, 278–287.
- Kim, P. R.; Carlson, K. E.; Katzenellenbogen, J. A. Progesterin 16 alpha-17-alpha-dioxolane ketals as molecular probes for the progesterone receptor: Synthesis, binding affinity, and photochemical evaluation. *J. Med. Chem.* **1993**, *36*, 1111–1119.
- Vijayaraghavan, S. T.; Balasubramanian, T. R. Synthesis of 3,4-diaryl-2-(5H)furanones. *Ind. J. Chem.* **1986**, *25B*, 760–761.
- Khanna, I. K.; Wei, R. M.; Yu, Y.; Xu, X. D.; Koszyk, F. J.; Collins, P. W.; Koboldt, C. M.; Veenhuizen, A. W.; Perkins, W. E.; Casler, J. J.; Masferrer, J. L.; Zhang, Y. Y.; Gregory, S. A.; Seibert, K.; Isakson, P. C. 1,2-Diarylimidazoles as potent, cyclooxygenase-2 selective, and orally active antiinflammatory agents. *J. Med. Chem.* **1997**, *40*, 1634–1647.
- Li, J. J.; Anderson, G. D.; Burton, E. G.; Nita Cogburn, J.; Collins, J. T.; Garland, D. J.; Gregory, S. A.; Huang, H.-C.; Isakson, P. C.; Koboldt, C. M.; Logusch, E. W.; Norton, M. B.; Perkins, W. E.; Reinhard, E. J.; Seibert, K.; Veenhuizen, A. W.; Zhang, Y.; Reitz, D. B. 1,2-Diarylcyclopentenones as selective cyclooxygenase-2 inhibitors and orally active antiinflammatory agents. *J. Med. Chem.* **1995**, *38*, 4570–4578.
- Lieber, E.; Curtice, J.; Rao, C. Structure and reactivity of organic azides. *Chem. Ind.* **1966**, 586.
- Pauling, L. In *The Nature of the Chemical Bond*, 3rd ed.; Cornell University Press: Ithaca, New York, 1960.
- Livingston, R.; Ramachandra Rao, C. An electron diffraction investigation of the molecular structure of methyl azide. *J. Phys. Chem.* **1960**, *64*, 756–759.
- Hansch, C.; Leo, A.; Unger, S. H.; Kim, K. H.; Nikaitani, D.; Lien, E. J. Aromatic substituent constants for structure activity correlations. *J. Med. Chem.* **1973**, *16*, 1207–1216.
- Sahi, S.; Srinivasan, M.; Kothekar, V. 530 ps molecular dynamics simulation of indoprofen and NS-398 with COX-1 and COX-2. Study of perturbative changes in the complexes. *J. Mol. Struct. (THEOCHEM)* **2000**, *498*, 133–148.
- Talley, J. J. Selective inhibitors of cyclooxygenase-2 (COX-2). *Prog. Med. Chem.* **1999**, *36*, 201–234.
- Chan, C.-C.; et al. Rofecoxib [Vioxx, MK-0966; 4-(4'-Methylsulfonylphenyl)-3-phenyl-2-(5H)-furanone]: A potent and orally active cyclooxygenase-2 inhibitor. Pharmacological and biochemical profiles. *J. Pharmacol. Exp. Ther.* **1999**, *290*, 551–560.
- Kumar, P.; Knaus, E. E. Synthesis and antiinflammatory activity of N-substituted-dihydropyridylacetic acids, esters and amides. *Drug Des. Delivery* **1987**, *2*, 145–149.
- Buolamwini, J. K.; Knaus, E. E. Synthesis and antinociceptive activity of 4-pyridyl and -dihydropyridyl analogues of meperidine and ketobemidone. *Drug Des. Delivery* **1990**, *7*, 19–31.